

# Theoretical Insight into the Structure, Energetic Property and Thermal Stability of $C_6N_6H_{12}$ Cages

Zhang J<sup>1\*</sup>, Chen G<sup>1</sup> and Gong X<sup>2</sup>

<sup>1</sup>College of Material and Chemical Engineering, ChuZhou University, ChuZhou, Anhui, PR China

<sup>2</sup>School of Chemical Engineering, Nanjing University of Science and Technology, Nanjing, Jiangsu, PR China

## Abstract

Recent theoretical studies have identified carbon-nitrogen cages are potentially stable high energy density materials. Two such  $C_6N_6H_{12}$  cages were proposed and investigated using the density functional theory comparison with two similar common cage compounds hexaazaisowurtzitane and cubane. The structure and energetic property were researched. Stability is approached in two ways: (1) stability of one isomer versus another based on the relative energy, (2) thermal stability determined by the bond breaking energies. Taking into consideration of the stability and detonation performance, two  $C_6N_6H_{12}$  cages may be candidate as potential high energy density compounds.

**Keywords:** Cage; Heat of formation; Strain energy; Detonation performance; Stability

## Introduction



Nitrogen molecules have been the research focus in recent years as possible candidates of high energy density compounds (HEDC). A molecule  $N_x$  can undergo the reaction  $N_x \rightarrow (x/2) N_2$ , a reaction that can be exothermic by more than 50 kcal/mol or more per nitrogen atom [1,2]. All-nitrogen molecules i.e.,  $N_4$ ,  $N_5$ ,  $N_6$ ,  $N_7$ ,  $N_8$ ,  $N_{10}$ ,  $N_{12}$ ,  $N_{18}$ ,  $N_{20}$ ,  $N_{24}$ ,  $N_{30}$ , and  $N_{36}$  have been studied theoretically [2-21]. There is also some research on nitrogen-rich molecules such as  $C_xN_y$  and  $O_xN_y$  species [22-26]. However, the properties of this kind of compounds are not very perfect, such as rare existence in experiments and not high stability. It has been shown that even the most stable  $N_{12}$  cage is of instability with regard to dissociation [27-29]. Thus, chemists often stabilize these all-nitrogen molecules with different atoms partially replacement of nitrogen atoms by sulfur atom, carbon atom or by oxygen atoms [16,17,22-26,30-32]. The advantages of these molecules are that they tend to have significant degrees of strain and high nitrogen content which will result in larger heat of formation and thus greater heat release in the detonation reactions as HEDCs. It is this point of view prompted us to research the analogy. In fact, one kind of similar  $C_5N_5H_{11}$  cages have been investigated in recent work [26]. In the current study, two  $C_6N_6H_{12}$  cages (Figure 1) with high symmetry and ring skeleton modified based on the most stable  $N_{12}$  cage are proposed and investigated comparison with two similar cage compounds hexaazaisowurtzitane (IW) and cubane. The structure, thermal stability, and energetic property including thermodynamic property, strain energy, heat of formation, and so on were studied through theoretical calculation for screening potential HEDCs. It is expected that our results could provide some novel insight for design and development of new HEDCs.

## Computational Details

Gaussian 03 program package [33] was used for all calculations in this study. The geometries of  $C_6N_6H_{12}$  cages are optimized with the density functional theory (DFT) at the B3LYP level along with 6-31+G(d,p) basis set [34-36]. Vibrational analysis is performed with the absence of imaginary vibrational frequency confirming the obtained structures correspond to the true minimum on the potential energy hypersurface. Ring strain energy (SE) is a significant parameter for cage compounds. It is defined as an energy excess through a reaction in which the cyclic molecule as reactant and a strain-free counterpart

as product. Various reactions have been suggested in the literature ranging from isogyric to isodesmic to homodesmotic forms [37,38]. Among them, the relatively accurate is homodesmotic schemes [39-43] in which not only the number of various bonds is conserved but also the valence environment around each atom is preserved to raise the advantage of canceling of systematic errors [44].

In the present work, the homodesmotic reactions designed for calculating the strain energies of S1 and S2 are as follows, respectively:



The change in energy, with the correction of zero-point vibrational energy (ZPE) of reactions is defined as the SE:

$$\Delta E = \sum E_0(\text{product}) - \sum E_0(\text{reactant}) + \Delta ZPE \quad (3)$$

Kamlet and Jacobs provided important insight into factors that determine detonation performance [45] measured by detonation velocity and detonation pressure which is expressed by the following equation:

$$D(\text{km} \cdot \text{s}^{-1}) = 1.01 [N^{0.5} M^{0.25} Q^{0.25} (1 + 1.30\rho)] \quad (4)$$

$$P(\text{Gpa}) = 1.558 \rho^2 N M^{0.5} Q^{0.5} \quad (5)$$

where  $D$  is detonation velocity,  $P$  is detonation pressure,  $\rho$  is the density of explosive in  $\text{g} \cdot \text{cm}^{-3}$ .  $N$  is the moles of gaseous detonation products

\*Corresponding author: Zhang J, College of Material and Chemical Engineering, ChuZhou University, ChuZhou, 239000, Anhui, PR China, Tel: +8615755054682; E-mail: zhangjy13@163.com

Received October 19, 2016; Accepted October 24, 2016; Published October 30, 2016

Citation: Zhang J, Chen G, Gong X (2016) Theoretical Insight into the Structure, Energetic Property and Thermal Stability of  $C_6N_6H_{12}$  Cages. J Phys Chem Biophys 6: 228. doi:10.4172/2161-0398.1000228

Copyright: © 2016 Zhang J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

per gram of explosives,  $M$  is the molecular weight of the detonation products, and  $Q$  is the heat release during the detonation reaction in cal·g<sup>-1</sup>.  $N$ ,  $M$ ,  $Q$  are determined based on the mot exothermic principle, that is the final detonation products are assumed N<sub>2</sub>(g), H<sub>2</sub>O(g), CO<sub>2</sub>(g), and C(g), with oxygen being used to form H<sub>2</sub>O before CO<sub>2</sub> for C, H, O, N-containing explosives.

Pyrolysis mechanism and thermal stability can be evaluated using bond dissociation energy (BDE) [20-25,46-51] which is a measure of the strength of a chemical bond. It can be defined as the enthalpy change at 298K and 1 atm in a molecule as follows:



$$BDE(R-X)=[HOF(R\cdot)+HOF(X\cdot)]-HOF(RX) \quad (7)$$

where,  $RX$  represents the neutral molecules,  $R\cdot$  and  $X\cdot$  represent the corresponding radical products produced by breaking the  $R-X$  bond.  $BDE(R-X)$  is the bond dissociation energy of bond  $R-X$ ,  $HOF(X\cdot)$ , and  $HOF(RX)$  are the standard heats of formation for the products and reactant at 298K, respectively. However, the BDE in the present work is defined as the difference between the total energies at 0 K in the unimolecular bond dissociations. Therefore, we computed the BDE at 0 K according to the energy changes during the bond process at the B3LYP/6-31G+ (d,p) level.

$$BDE(RX)=\Delta E=[E(R\cdot)+E(X\cdot)-E(RX)] \quad (8)$$

## Results and Discussion

### Molecular structure and electronic property

The C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cage of S2 is modified based on the most stable N<sub>12</sub> cage. It has four pentagons and two hexagons while S1 is composed of five hexagons as the derivative of S2. Table 1 shows the representative parameters with hexaazaisowurtzitane (IW) for comparison. It can be seen that all the C-C bonds of two C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cages and IW are longer than the normal C-C bond (1.540Å), except for C1-C6 and C7-C8 bonds in S2, and the largest one reaches 1.611 Å for C1-C9 bond which is between two six-membered rings in S1, C1-C6 bond 1.595 Å between two heptagons in IW. Meanwhile, the C1-C6 and C7-C8 bonds in S2 which located at the junction of two five-membered rings and C3-C4 and C8-C9 bonds in IW between pentagon and hexagon are shorter than the normal C-C bond (1.540Å).

Meanwhile, all the bond lengths of C-N bonds in S1 are within normal range (1.470 Å), while C-N bonds in S2 are other different case. The longest C-N bond is C7-N11 (1.538 Å) in S2 which located at the junction of pentagon and hexagon. The smallest bond angles in S1, S2, and IW are 100.2, 98.7, and 102.3°, respectively, which are smaller than the usual bonding angles of tetrahedral carbon (109.47°) and trivalent nitrogen (107.30°) but larger than 90° in cubane. It can be construed to mean that these cage structures subject to a certain degree of strain but weaker than that in cubane and they be released as an additional energy during the detonation.

Table 2 collects the frontier molecular orbital energies, the highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO) energies and the energy gaps ( $\Delta E_{LUMO-HOMO}$ ) for S1 and S2. The energy gap  $\Delta E_{LUMO-HOMO}$  is an important indicator of stability for molecules. The larger the  $\Delta E_{LUMO-HOMO}$  is, the more stable the molecule is during the chemical or photochemical process with electron transfer or electron leap. For example, S1 has a smaller  $\Delta E_{LUMO-HOMO}$  than that of S2, indicating that S1 is less stable than that of S2 when a chemical or photochemical reaction happens.

### Infrared spectrum and thermodynamic property

Infrared spectrum (IR) is not only the basic property of compounds, but also an efficient approach to identify substances. Besides, it is necessary to predict the thermodynamic properties. Therefore, it is a very important parameter. Since the DFT-calculated harmonic vibrational frequencies are usually larger than those observed experimentally, they are scaled by using a factor of 0.96 [52] which has been proved to be suitable [21-25,48,50]. Figure 2 presents the simulated IR spectrum for S1 and S2. Obviously, there are four main characteristic regions in S1. The modes in 574.57 and 859.21 cm<sup>-1</sup> are associated with the C-H twisting out of plane. The peak at 713.15 cm<sup>-1</sup> is characterized to the N-H twisting out of plane. The band at 3070.77~ 3087.88 cm<sup>-1</sup> is associated with the C-H stretch. Meanwhile, there are three main characteristic regions in S2. The strongest modes in 815.07 cm<sup>-1</sup> are associated with the N-H rocking vibration. The band at 3092.17~ 3151.09 cm<sup>-1</sup> is associated with the C-H stretch. Also, there are many small peaks in S2, which can be attributable to its skeleton distortions caused by many modes of vibration from chemical bonds.

Thermodynamic properties are important parameters, i.e., standard molar heat capacity  $C_{p,m}^{\circ}$  in J·mol<sup>-1</sup>·K<sup>-1</sup>, entropy  $S_m^{\circ}$  in J·mol<sup>-1</sup>·K<sup>-1</sup>, and enthalpy  $H_m^{\circ}$  in kJ·mol<sup>-1</sup>. Based on the scaled vibrational frequencies,  $C_{p,m}^{\circ}$ ,  $S_m^{\circ}$ , and  $H_m^{\circ}$  of S1 and S2 ranging from 200 to 800 K are obtained based on statistical thermodynamic principle and listed in Table 3. It can be seen that all the thermodynamic properties increase with the temperature evidently. This is because the main contributions to the thermodynamic functions are from the translation and rotation of molecules when the temperature is low; however, at higher temperature, the vibrations are intensified and therefore make more contributions to the thermodynamic properties and lead to the increase in the thermodynamic functions. In addition, the  $C_{p,m}^{\circ}$ ,  $S_m^{\circ}$ , and  $H_m^{\circ}$  of S1 are generally larger than those of S2. This could be due to the fact that S1 has one more C-C bond than that of S2 while they have the same numbers of C-N bonds, resulting in more freedoms of motion in S1, i.e., translation, rotation, and vibration, thus more contributions to the thermodynamic properties for S1. The temperature-dependent relations for  $C_{p,m}^{\circ}$ ,  $S_m^{\circ}$ , and  $H_m^{\circ}$  are as follows and expressed as in Figure 3.

$$S1: C_{p,m}^{\circ} = -89.91 + 1.01T - 7.06 \times 10^{-4}T^2, R^2=0.9997, SD=1.4907 \quad (9)$$

$$S_m^{\circ} = 184.08 + 0.58T - 2.39 \times 10^{-4}T^2, R^2=0.9995, SD=0.1516 \quad (10)$$

$$H_m^{\circ} = -6.37 + 0.02T + 2.53 \times 10^{-4}T^2, R^2=0.9995, SD=0.1865 \quad (11)$$

$$S2: C_{p,m}^{\circ} = -87.80 + 0.95T - 4.73 \times 10^{-4}T^2, R^2=0.9997, SD=1.6726 \quad (12)$$

$$S_m^{\circ} = 191.93 + 0.52T - 1.03 \times 10^{-4}T^2, R^2=0.9994, SD=2.0877 \quad (13)$$

$$H_m^{\circ} = -4.82 + 0.02T + 2.38 \times 10^{-4}T^2, R^2=0.9995, SD=0.3004 \quad (14)$$

### Strain energy and heat of formation

The assessment of ring strain energy in ring compounds has been a topic of major interest to the organic chemist for decades [49-58]. Studies have proved that computational results of SE at the B3LYP/6-31G+(d,p) via homodesmotic process are reasonable [22,23,37,38,59]. For example, the SE of norbornane obtained by the homodesmotic method (63.15 kJ·mol<sup>-1</sup>) [60] is in agreement with experimental value (60.22 kJ·mol<sup>-1</sup>) [61]. In the present work, the strain energies of C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cages are obtained by the homodesmotic reaction (1) and (2). For comparison, the following reactions are used to evaluate the SEs of hexaazaisowurtzitane and cubane, respectively:




Figure 1: The structures of two  $C_6N_6H_{12}$  cages, hexaazaisowurtzitane, and cubane (H atoms omitted).




Figure 2: The calculated IR of  $C_6N_6H_{12}$  cages.

S1		S2		IW	
C1-C9 1.611	C9-N12 1.468	C1-C6 1.518	C8-N2 1.449	C1-C6 1.595	C4-N12 1.449
C2-C8 1.610	C7-N10 1.468	C7-C8 1.518	C8-N10 1.530	C3-C4 1.575	C6-N7 1.466
C4-C7 1.610	C8-N12 1.468	C1-N3 1.449	C9-N4 1.491	C8-C9 1.575	C6-N10 1.472
C1-N5 1.468	C4-N3 1.468	C1-N10 1.530	C9-N5 1.491	C1-N2 1.472	C8-N7 1.496
C1-N6 1.468	C2-N3 1.468	C6-N4 1.442	C9-N10 1.531	C1-N5 1.466	C8-N11 1.449
C9-N10 1.468		C6-N11 1.538	C12-N2 1.458	C3-N2 1.481	C9-N10 1.481
		C7-N5 1.442	C12-N3 1.458	C3-N11 1.462	C9-N12 1.462
		C7-N11 1.538	C12-N11 1.584	C4-N5 1.584	
N3-C4-N5 105.7	N5-C4-N3 105.7	C1-N3-C12 100.6	N4-C6-N11 118.8	C1-N2-C3 105.2	C3-C4-N5 105.6
N3-C2-N6 105.7	N5-C4-C7 111.8	C1-C6-N4 99.4	C6-C1-N3 104.3	C1-N5-C4 103.8	C3-C4-N12 109.2
C2-N3-C4 108.4	C4-C7-N11 111.7	C1-C6-N11 98.9	C6-C1-N10 98.7	C1-C6-N7 110.1	C4-N12-C9 113.5
C2-N6-C1 108.5	C7-N11-C8 100.2	N2-C8-C7 104.3	C7-N5-N9 100.9	C1-C6-N10 114.6	C8-N11-C3 113.5
N6-C1-N5 105.7	N11-C8-N12 105.7	N2-C8-N10 114.3	N5-C9-N4 108.2	N2-C3-C4 102.3	N7-C8-N11 113.1
C1-N5-C4 108.5	C8-N11-C9 108.5	N2-C12-N3 102.3	N5-C9-N10 105.6	N2-C3-N11 116.4	C8-C9-N12 109.8

Table 1: Bond lengths (Å) and bond angles ( $^{\circ}$ ) of two  $C_6N_6H_{12}$  cages and IW.






Table 4 collects the total energies ( $E_0$ ) and SEs of the species involved in the reactions. Obviously, S2 has a larger strain energy ( $3475.88 \text{ kJ}\cdot\text{mol}^{-1}$ ) than that of S1 ( $287.93 \text{ kJ}\cdot\text{mol}^{-1}$ ), which can be attributable to its special structures composed of four pentagons and two hexagons while S1 is made up of five hexagons. As is known, the

smaller the ring is, the greater distorted the molecule is, thus, the less the strain energy becomes. Thus, it could be concluded that S1 is more stable than S2 from this point of view.

A key property of an energetic material that is used to assess its potential performance in a gun or warhead is its heat of formation (HOF), as it enters into the calculation of explosive and propellant properties such as detonation velocity and detonation pressure. DFT methods have been proved accurate for computing HOF through appropriate reactions [21-25,62-65]. In the present work, we calculated the HOFs of S1 and S2 by reaction (1) and (2) and hexaazaisowurtzitane by reaction (17) as well as the HOF of cubane by reaction (18) for comparison.



Isomers	$E_{\text{HOMO}}/\text{eV}$	$E_{\text{LUMO}}/\text{eV}$	$\Delta E_{\text{LUMO-HOMO}}/\text{eV}$
S1	-0.18922	-0.01954	0.16968
S2	-0.23056	-0.00926	0.22130

Table 2: Selected properties of two C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cages.



Compd.	Parameters	200.0K	300.0K	400.0K	500.0K	600.0K	700.0K	800.0K
S1	$C_{p,m}^0$	92.75	165.82	234.65	290.61	334.22	368.51	396.11
	$S_m^0$	301.80	353.05	410.44	469.06	526.06	580.26	631.33
	$H_m^0$	9.74	22.63	42.74	69.12	100.45	135.65	173.93
S2	$C_{p,m}^0$	84.06	150.38	216.22	269.34	310.46	342.49	368.02
	$S_m^0$	298.89	345.32	397.92	452.12	505.02	555.38	602.84
	$H_m^0$	9.37	21.05	39.48	63.87	92.95	125.66	161.24

Table 3: Thermodynamic properties of S1 and S2 at different temperatures.

Compd.	$E_0^a$ (a.u.)	ZPE/(a.u.)	SE/(kJ·mol <sup>-1</sup> )	HOF/(kJ·mol <sup>-1</sup> )
CH <sub>3</sub> NHCH <sub>3</sub>	-135.18128	0.092350	0	-82.98 <sup>c</sup>
NH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	-151.23042	0.082069	0	-78.84 <sup>c</sup>
CH <sub>3</sub> NH <sub>2</sub>	-95.87185	0.064051	0	-80.37 <sup>c</sup>
CH <sub>3</sub> CH <sub>3</sub>	-79.84164	0.074599	0	-84.00 <sup>d</sup>
S1	-564.32847	0.21048	287.93	520.85
S2	-563.09026	0.18642	3475.88	3711.86
IW	-564.36584	0.21129	192.16	425.08 (319.88 <sup>e</sup> )
cubane	-309.47990	0.13339	573.91 (656.57 <sup>e</sup> )	507.91 (619.98 <sup>e</sup> )

<sup>a</sup> $E_0$  is the total energy after the correction of ZPE; <sup>b</sup>Ref [62]; <sup>c</sup>Obtained at the G3 level from the formation reactions: 2C(g)+3H<sub>2</sub>+N<sub>2</sub>→CH<sub>3</sub>NHCH<sub>3</sub>, C(g)+3H<sub>2</sub>+N<sub>2</sub>→NH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, C(g)+2.5H<sub>2</sub>+0.5N<sub>2</sub>→CH<sub>3</sub>NH<sub>2</sub>; <sup>d</sup>Ref [63]; <sup>e</sup>Ref [37]; <sup>f</sup>Ref [64]

Table 4: SEs and HOFs for C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cages, IW, and cubane.



From the data in Table 4, it can be seen that the strain energy of S2 is seven times larger than that of S1 and both of them are not only larger than those of IW either obtained at the same computational level (425.08 kJ·mol<sup>-1</sup>) or experimental value (319.88 kJ·mol<sup>-1</sup>) but also larger than that of cubane at the same computational level (507.91 kJ·mol<sup>-1</sup>). This indicates that the heat release will be gained in both of C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cage and S2 will benefit more during the detonation.

### Density and detonation characteristics


A high density is a top priority in seeking to develop new explosives. Then follow the detonation velocity and detonation pressure which are also crucial importance to an energetic material. In the present work, density was predicted from the molecular volume divided by molecular weight, while the molecular volume was obtained from the average of 100 single points molar volumes for each optimized structures. Detonation velocity  $D$  and detonation pressure  $P$  are evaluated by the Kamlet and Jacobs equation (4) and (5).

The amount of energy  $Q$ , important parameters reflecting the explosive performances of energetic materials, is commonly taken to corresponding to the total energy change per gram during the detonation. It is the negative of the total energy change:

$$Q = -\frac{1}{M} \left[ \sum_p n_p \text{HOF}(p) - \text{HOF}(A) \right] \quad (19)$$

Where  $n_p$  is the number of moles of final product  $p$ , having molar heat of formation HOF( $p$ ), and HOF( $A$ ) is the molar heat of formation of the compound  $A$ .

The following detonation reactions were used to evaluate the detonation energy  $Q$  for two C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> isomers, IW, and cubane:








Table 5 collects the densities and detonation characteristics for the corresponding compounds. It is noted that both of S1 and S2 have larger density than that of cubane regardless of computational result (1.235 g·cm<sup>-3</sup>) obtained at the same level or experimental value (1.290 g·cm<sup>-3</sup>). The detonation velocity of S1 is slight smaller than that of S2 but still larger than that of cubane. Both of S1 and S2 have higher detonation pressure that those of IW and cubane indicating that S1 and S2 have greater power during the detonation process. The detonation energy of S1 and S2 are more negative than those of IW and cubane, implying that the title cages will release more energy during the detonation. It might seem surprising that the Q of S2 is rather negative which may be caused by its high HOF and large SE.

### Thermal stability

Another important aspect of energetic materials is whether they are kinetically stable enough to be of practical interest. A good candidate HEDM should have a high stability. Thus, studies on the thermal stability are important and essential for understanding the decomposition process of the energetic materials because they are directly relevant to the sensitivity and safety of the energetic compounds during the storage or transportation.

In the present work, the stability of C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cages is approached in two ways: (1) stability of one isomer versus another based on the relative energy, (2) thermal stability determined by the bond breaking energies. And so we can get a better understanding of the relationship between the structures and thermal stability for this kind of cage compounds.

Table 6 collects the relative energies of two C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cages obtained at the B3LYP/6-31G+(d,p) level and their composition. Obviously, S1 gets a more negative energy than that of S2, thus, S1 is more stable than S2. The primary reason may be the relative bond strength between C-C and C-N bonds. The bond energies for two bonds are 347.94 and 304.87 kJ·mol<sup>-1</sup>, respectively [66-71]. Generally, for any rearrangement of the atoms, the replacement of a C-C with C-N bond should be energetically disadvantageous by 43.07 kJ·mol<sup>-1</sup>. Compared with S2, S1 should reap the benefit for its arrangement and should be more stable by about 652.81 kJ·mol<sup>-1</sup>. Another reason may be due to their structures. S1 is composed by five hexagons while S2 is made up of four pentagons and two hexagons. As is known, the smaller the polygon is, the larger SE it subjects to. Therefore, S2 experiences more strain energy which has been verified in the section of “strain energy” and it is less stable than S1.

The bond breaking energy is another important factor that related to the stability for energetic materials. The bond strength of C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cages is evaluated by the Wiberg Bond Index (WBI) based on the NBO analysis. Often, the smaller Wiberg bond index is, the weaker bond is. Therefore, we picked out the smallest WBI of the same kind.

For example, the environments of C1-C9, C2-C8, and C4-C7 in S2 consist of four C-N bonds and both of them are at the junction of two hexagons, that is, they belong to the same kind of bond. In the same way, we select C1-C6, C1-N3, C1-N10, C9-N4, and C12-N11 bonds for S2, C1-C6 and C4-N5 bonds for IW, and C1-C3 bond for cubane.

From the results in Table 7, one can find the weakest bonds for S1, S2, IW, and cubane are C2-N3, C1-N10, C1-C6, and C1-C3 with BDEs 240.14 kJ·mol<sup>-1</sup>, 134.49 kJ·mol<sup>-1</sup>, 263.67 kJ·mol<sup>-1</sup>, and 160.46 kJ·mol<sup>-1</sup>, respectively, indicating that S1, S2, and IW will break a C-N bond for the first step in the pyrolysis while it would begin with the breaking of a C-C bond for cubane. In addition, S1 has a better stability than S2 although both of them may not as stable as IW. In addition, the results in the present work also prove that the dissociation energy of C-C bond is not always larger than that of C-N bond in the cage molecules.

### Conclusions

On the basis of theoretical studies on two C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cages comparison with hexaazaisowurtzitan and cubane, we can draw the following conclusions:

- (1) The thermodynamic properties of two cages all increase with the temperature remarkably. S1 has generally larger C<sub>p,m</sub><sup>o</sup>, S<sub>m</sub><sup>o</sup>, and H<sub>m</sub><sup>o</sup> than those of S2 at the corresponding temperature due to their different structure composition.
- (2) The results of strain energies indicate S2 composed of more small polygons experiences much larger strain energy than S1 which may lead to the instability of S2.
- (3) Both of S1 and S2 have high positive heat of formation. Thus, quite huge of energy will be released if the compounds decomposes which verifies the heat release during the detonation process.
- (4) The density and detonation characteristics of S1 and S2 are equivalent or slightly superior to those of hexaazaisowurtzitan and cubane, especially they have relatively negative detonation

Compd.	$\rho$ (g·cm <sup>-3</sup> )	D/(km·s <sup>-1</sup> )	P/(Gpa)	Q/(kJ·g <sup>-1</sup> )
S1	1.418	5.72	12.43	-25.79
S2	1.498	5.93	13.88	-43.57
IW	1.441 (1.570 <sup>a</sup> )	5.78 (6.12 <sup>a</sup> )	12.84 (15.24 <sup>a</sup> )	-25.22
cubane	1.235 (1.290 <sup>b</sup> )	3.59 (3.69 <sup>b</sup> )	4.42 (4.82 <sup>b</sup> )	-21.20

<sup>a</sup>Calculated value from Ref. [65]; <sup>b</sup>Experimental value from Ref [64].

**Table 5:** Densities and Enthalpies of combustion for C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> cages, IW, and cubane.

Isomers		S1	S2
Bonds	C-C bonds	3	2
	C-N bonds	12	11
Polygons	pentagons	0	4
	Hexagons	5	2
Energies (kJ·mol <sup>-1</sup> )		-774.73	0

**Table 6:** Bond properties and relative energies of two cages.

	C-C		C-N		
	Bond	WBI	Bond	BDE	
S1	C1-C9	0.9203	C2-N3	0.9883	
		323.12	C1-N3	0.9991	
S2	C1-C6	0.9671	170.27	C1-N10	0.9350
				C9-N4	0.9710
				C12-N11	0.8981
				C4-N5	0.9585
IW	C1-C6	0.9266	263.27		
Cubane	C1-C3	0.9917	160.46		

**Table 7:** WBI and BDE (kJ·mol<sup>-1</sup>) for C<sub>6</sub>N<sub>6</sub>H<sub>12</sub> isomers, IW, and cubane.

energy. (5) The results of stability suggest S1 which contains more C-C bonds are usually more stable than S2 in which C-N bonds are more than C-C bonds and also S1 is more stable with respect to bond dissociation energy. Taking the detonation performance and stability into consideration, two  $C_6N_6H_{12}$  cages may be candidate as potential HEDMs. It is hoped that the present systemic comparative investigation would be useful for molecular structural design and research of cage compounds in future.

#### Acknowledgements

This project is supported by the research fund of ChuZhou University under No. 2014qd039, 2015qd13, and General project of natural science research in Anhui Province (KJ2015B08).

#### References

- Lauderdale WJ, Stanton JF, Barlett RJ (1992) Stability and energetics of metastable molecules: tetraazatetrahedrane ( $N_4$ ), hexaazabenzene ( $N_6$ ), and octaazacubane ( $N_8$ ). *J Phys Chem* 96: 1173-1178.
- Glukhovtsev MN, Jiao H, Schleyer PV (1996) Besides  $N_2$ , What Is the Most Stable Molecule Composed Only of Nitrogen Atoms?. *Inorg Chem* 35: 7124-7133.
- Leininger ML, Van Huis TJ, Henry III FS (1997) Protonated High Energy Density Materials:  $N_4$  Tetrahedron and  $N_8$  Octahedron. *J Phys Chem A* 101: 4460-4464.
- Zheng JP, Jacek W, Jens SL, Daniel MB, Radziszewski JG (2000) Tetrazete ( $N_4$ ). Can it be prepared and observed? *Chem Phys Lett* 328: 227-233.
- Fau S, Wilson K J, Bartlett RJ (2002) On the stability of  $N_5^{+}N_5^{-}$ . *J Phys Chem A* 106: 4639-4644.
- Gagliardi L, Evangelisti S, Vincenzo B, Björn OR (2000) On the dissociation of  $N_8$  into  $3N_2$  molecules. *Chem Phys Lett* 320: 518-522.
- Wang LJ, Peter W, Paul GM (2002) Theoretical Prediction on the Synthesis Reaction Pathway of  $N_6$  ( $C_{2h}$ ). *J Phys Chem A* 106: 2748-2752.
- Ray E (1992) Ab-Initio Correlated Calculations of six  $N_2$  Isomers ( $N_8$ ). *J Phys Chem* 96: 10789-10792.
- Motoi T, Rodney J B (2001) Structure and Stability of  $N_6$  Isomers and Their Spectroscopic Characteristics. *J Phys Chem A* 105: 4107-4113.
- Li QS, Hu XG, Xu WG (1998) Structure and stability of  $N_7$  cluster. *Chem Phys Lett* 287: 94-99.
- Engelke R, Stine JR (1990) Is  $N_8$  cubane stable?. *J Phys Chem* 94: 5689-5694.
- Schmidt MW, Gordon MS, Boatz JA (2000) Cubic fuels?. *Int J Quantum Chem* 76: 434-446.
- Leininger ML, Sherrill CD, Schaefer HF (1995)  $N_8$ : A Structure Analogous to Pentalene, and Other High Energy Density Minima. *J Phys Chem* 99: 2324-2328.
- Tian A, Ding F, Zhang L, Xie Y, Schaefer HF (1997) New Isomers of  $N_8$  without Double Bonds. *J Phys Chem A* 101: 1946-1950.
- Strout DL (2002) Acyclic  $N_{10}$  fails as a high energy density material. *J Phys Chem A* 106: 816-818.
- Brune LY, Bledson TM, Strout DL (2003) What makes an  $N_{12}$  cage stable?. *Inorg Chem* 42: 8117-8120.
- Sturdivant SE, Nelson FA, Strout DL (2004) Trends in Stability for  $N_{18}$  Cages. *J Phys Chem A* 108: 7087-7090.
- Bliznyuk AA, Shen M, Schaefer HF (1992) The dodecahedral  $N_{20}$  molecule: Some theoretical predictions. *Chem Phys Lett* 198: 249-252.
- Ha TK, Suleimenov O, Nguyen MT (1999) A quantum chemical study of three isomers of  $N_{20}$ . *Chem. Phys Lett* 315: 327-334.
- Strout DL (2005) Why Isn't the  $N_{20}$  Dodecahedron Ideal for Three-Coordinate Nitrogen?. *J Phys Chem A* 109: 1478-1480.
- Strout DL (2004) Isomer Stability of  $N_{24}$ ,  $N_{30}$ , and  $N_{36}$  Cages: Cylindrical versus Spherical Structure. *J Phys Chem A* 108: 2555-2558.
- Wang LJ, Zgierski MZ, Mezey PG (2003) Stable Structures of Nitrogen-Rich Sulfides:  $S(N_3)_m$  ( $m = 1-4$ ). *J Phys Chem A* 107: 2080-2084.
- Cheng LP, Li QS (2004) *J Phys Chem A* 108: 165.
- Brune LY, Strout DL (2003) Are Four Oxygens Enough to Stabilize the  $N_8$  Cube? Dissociation Pathways of Cage Isomers of  $N_8O_4$ . *J Phys Chem A* 107: 5840-5843.
- Strout DL (2003) Stabilization of an all-nitrogen molecule by oxygen insertion: dissociation pathways of  $N_8O_6$ . *J Phys Chem A* 107: 1647-1650.
- Zhang JY, Gong XD (2015) Comparative theoretical investigation of the structures, energetics, and stabilities of  $C_7N_5H_{11}$  cages *J Mol Model* 21: 1-8.
- Li QS, Zhao JF (2002) Theoretical Study of Potential Energy Surfaces for  $N_{12}$  Clusters. *J Phys Chem A* 106: 5367-5372.
- Douglas LS (2006) Isomer stability of  $N_6C_6H_6$  Cages. *J Phys Chem A* 110: 7228-7231.
- Karleta DC, Roshawnda C, Douglas LS (2006) Stability of Carbon-Nitrogen Cages in 3-Fold Symmetry. *J Chem Theory Comput* 2: 25-29.
- Jursic BS (2000) Structural, energetic, and vibrational features of azacubanes as energetic materials studied with density functional theory methods. *J Mol Struct* 530: 21-30.
- Sturdivant SE, Strout D (2004) Beyond  $N_8O_6$ : Length effects and end-cap effects on the stability of a hexagonal nitrogen tube. *J Phys Chem A* 108: 4773-4777.
- Colvin KD, Strout D (2005) Stability of nitrogen-oxygen cages  $N_{12}O_2$ ,  $N_{14}O_2$ ,  $N_{14}O_3$ , and  $N_{16}O_4$ . *J Phys Chem A* 109: 8011-8015.
- Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, et al. (2003) Gaussian 03, Revision A1; Gaussian, Inc.: Pittsburgh PA, USA.
- Becke AD (1993) Synthesis and structural studies of a new class of quaternary ammonium salts, which are derivatives of cage adamanzane type aminal 1, 3, 6, 8- tetraazatricyclo [4.3.1.13,8] undecane (TATU). *J Chem Phys* 98: 5648-5652.
- Becke AD (1988) Density-functional exchange-energy approximation with correct asymptotic behavior. *Phys Rev A* 38: 3098-3100.
- Hehre WJ, Radom L, Schleyer PR, Pople JA (1986) *Ab Initio Molecular Orbital Theory*. Wiley-Interscience: New York.
- George P, Trachtman M, Bock CW, Brett AM (1975) An alternative approach to the problem of assessing stabilization energies in cyclic conjugated hydrocarbons. *Theor Chim Acta* 38: 121-129.
- Zhu Y, Chen CX, Li XQ, Zhou Y, Huang W, et al. (2013) Association between blood pressure level and incidence of carotid artery plaque in middle-aged and elderly people. *Zhonghua Xin Xue Guan Bing Za Zhi* 41: 611-615.
- Zhao M, Gimarc BM (1993) strain energies in cyclic oxygen on,  $n = 3-8$ . *J Phys Chem* 97: 4023-4030.
- Bachrach SM (1989) Theoretical studies of phosphirane and phosphetane. *J Phys Chem* 93: 7780-7784.
- Magers DH, Davis SR (1999) Ring strain in the oxazetidines. *J Mol Struct* 487: 205-210.
- Yang J, Gong X, Wang G (2014) Structure, aromaticity, stability, and energetic performance of the analogues and derivatives of s-heptazine. *J Mol Model* 20: 1-10.
- Dill JD, Greenberg A, Liebman JF (1979) Substituent effects on strain energies. *J Am Chem Soc* 101: 6814-6826.
- Walker JE, Adamson PA, Davis SR (1999) Comparison of calculated hydrocarbon strain energies using ab initio and composite methods. *Comput Theor Chem* 487: 145-150.
- Kamlet MJ, Jacobs SJ (1968) Chemistry of detonations. I. A simple method for calculating detonation properties of C-H-N-O explosives. *J Chem Phys* 48: 23-35.
- Fan JF, Xiao HM (1996) Theoretical study on pyrolysis and sensitivity of energetic compounds. (2) Nitro derivatives of benzene. *J Mol Struct* 365: 225-229.
- Fan JF, Gu ZM, Xiao HM, Dong HS (1998) Theoretical study on pyrolysis and sensitivity of energetic compounds. (4) Nitro derivatives of phenols. *J Phys Org Chem* 11: 177-184.

48. Zhang J, Xiao HM (2002) Computational studies on the infrared vibrational spectra, thermodynamic properties, detonation properties and pyrolysis mechanism of octanitrocubane. J Chem Phys 116: 10674-10683.
49. Ju XH, Xiao HM, Xia QY (2003) A density functional theory investigation of 1,1-diamino-2,2-dinitroethylene dimers and crystal. J Chem Phys 119: 10247-10255.
50. Xu XJ, Xiao HM, Ju XH, Gong XD, Chen ZX (2005) Theoretical Studies on the Vibrational Spectra, Thermodynamic Properties, Detonation Properties and Pyrolysis Mechanisms for Polynitroadamantanes. J Phys Chem A 109: 11268-11274.
51. Qiu L, Xiao HM, Gong XD, Ju XH, Zhu WH (2006) Theoretical Studies on the Structures, Thermodynamic Properties, Detonation Properties, and Pyrolysis Mechanisms of Spiro Nitramines. J Phys Chem A 110: 3797-3807.
52. Scott P, Radom L (1996) Harmonic Vibrational Frequencies: An Evaluation of Hartree-Fock, Møller-Plesset, Quadratic Configuration Interaction, Density Functional Theory, and Semiempirical Scale Factors J Phys Chem 100: 16502-16513.
53. Alcamí M, Mo O, Yanez M (1998) G2 ab initio calculations on three-membered rings: Role of hydrogen atoms. J Comput Chem 19: 1072-1086.
54. Lewis LL, Turner LL, Salter EA, Magers DH (2002) Computation of the conventional strain energy in oxaziridine. J Mol Struct 592: 161-171.
55. Cremer D, Kraka E (1990) In Theoretical Models of the Chemical Bond. Maksia ZB (ed.), Springer, Heidelberg, Germany 2: 453.
56. Greenberg A, Liebman JF (2004) Strain energy of small ring hydrocarbons. Influence of CH bond dissociation energies. J Am Chem Soc 126: 4444-4452.
57. Bach RD, Dmitrenko O (2002) The effect of substituents on the strain energies of small ring compounds. J Org Chem 67: 2588-2599.
58. Bach RD, Dmitrenko O (2002) Effect of geminal substitution on the strain energy of dioxiranes. Origin of the low ring strain of dimethyldioxirane. J Org Chem 67: 3884-3896.
59. Wiberg KW, Fenoglio RA (1968) Heats of formation of C<sub>4</sub>H<sub>6</sub> hydrocarbons. J Am Chem Soc 90: 3395.
60. Dudev T, Lim C (1998) Ring strain energies from ab initio calculations. J Am Chem Soc 120: 4450-4458.
61. Bach RD, Dmitrenko O (2006) The effect of carbonyl substitution on the strain energy of small ring compounds and their six-member ring reference compounds. J Am Chem Soc 128: 4598-4611.
62. Zhao M, Gimarc BM (1993) Strain energies in cyclic oxygen On, n=3-8. J Phys Chem 97: 4023-4030.
63. Bachrach SM (1989) J Phys Chem 93: 7780.
64. Walker JE, Adamson PA, Davis SR (1999) Comparison of calculated hydrocarbon strain energies using ab initio and composite methods. Comput Theor Chem 487: 145-150.
65. Yao XQ, Hou XJ, Wu GS, Xu YY, Xiang HW, et al. (2002) Estimation of C-C Bond Dissociation Enthalpies of Large Aromatic Hydrocarbon Compounds Using DFT Methods. J Phys Chem A 106: 7184-7189.
66. Kybett BD, Carroll S, Natalis P, Bonnell DW, Margrove JL, et al. (1966) Thermodynamic properties of cubane. J Am Chem Soc 88: 626.
67. Lide DR (2010) CRC Handbook of Chemistry and Physics. 89th edn. CRC Press/Taylor and Francis, Boca Raton, FL, USA.
68. Slikder AK, Sikder N (2004) A review of advanced high performance, insensitive and thermal stable energetic materials emerging for military and space applications. J Hazard Mater 112: 1-15.
69. Liu H, Gong XD (2013) Comparative theoretical studies of substituted bridged bipyridines and their N-oxides. Struct Chem 24: 471-480.
70. Ghule VD, Jadhav PM, Patil RS, Radhakrishnan S, Soman T (2010) Quantum-chemical studies on hexaazaisowurtzitanes. J Phys Chem A 114: 498-503.
71. Atkins P, de Paula J (2002) Physical Chemistry. Bond Energies. 7th edn. Freeman WH and Co., New York, USA.