



Combining Ability of Inbred Lines of Maize (*Zea mays*) in Kiambu and Embu Counties, Kenya

J. M. Kariuki^{1*}, F. M. Njoka², P. K. Leley³ and D. W. Manene¹

¹Kenyatta University, Kenya.

²Embu University College, Kenya.

³KALRO Katumani Centre, Kenya.

Authors' contributions

This work was carried out in collaboration between all authors. Author JMK designed the study, wrote the protocol and wrote the first draft of the manuscript. Authors FMN and PKL were the research supervisors. Authors JMK and DWM read the first manuscript. Authors FMN and PKL managed the literature searches. Authors JMK and DWM managed the analyses of the study performed. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2016/27694

Editor(s):

(1) Fatemeh Nejatzaadeh, Department of Horticulture, Faculty of Agriculture, Khoy Branch, Islamic Azad University, Iran.

Reviewers:

(1) Abdulwahab Saliu Shaibu, Bayero University, Kano, Nigeria.

(2) Chunqing Zhang, Shandong Agriculture University, Taian, China.

(3) Gilberto Rodriguez Perez, Institute Technologic of Roque, Guanajuato, Mexico.

(4) Russel Pinnisch, Syngenta Seeds Inc, Janesville, USA.

Complete Peer review History: <http://www.sciencedomain.org/review-history/15981>

Original Research Article

Received 14th June 2016
Accepted 16th August 2016
Published 29th August 2016

ABSTRACT

Despite the fact that virtually all households in Kenya grow maize, over 60% of them are net maize buyers because they do not produce enough for their consumption. This is due to both biotic and abiotic factors such as poor planting materials, diseases and unreliable weather among others. This study was conducted to select lines with good combining ability. The trials were conducted in 2012 at experimental stations of Kenya Agricultural and Livestock Research Organization (KALRO) Muguga South and Embu in Kiambu and Embu counties of Kenya respectively. The study was conducted with 18 inbred lines. The experiment was laid out in a 6 x 6 lattice incomplete randomized block design with two replications. In Embu inbred line POPA produced the best grain yields when crossed with MUL 541 and MUL 521. Its high grain yield was also witnessed in Muguga where on average its performance was superior to other inbred lines. Inbred line MUL 513 can further be evaluated for grain yield improvement with all the other inbred lines which had high

*Corresponding author: E-mail: muthonicjeff@gmail.com;

grain yields. The best general combiners for grain yield were MUL 508, MUL 688, POP A, MUL 541, MUL 513 and MUL 114. POP A x MUL 541 produced the highest yield of 0.49 t/ha. Based on SCA estimates, the best cross combinations for plant height were MUL 508 x MUL 688, POP A x MUL541 and MUL513 x MUL114. For ear height best cross combinations were: MUL 508 x MUL 688, POP A x MUL 141, POP A x MUL 541 and MUL 513 x MUL 114. The best combinations for grain yield were MUL 508 x MUL 688, POP A x MUL 541 and MUL 513 x MUL 114. Crosses MUL508 x MUL 521, MUL 541 x MUL 508, and MUL 688 x MUL508 were good combinations for MSV disease resistance since in both research sites they had an MSV score of 1. For grain yield improvement on specific combining ability crosses MUL 508 x MUL 688, POP A x MUL 141, MUL 513 x MUL 114 and MUL 513 x CN244 can further be evaluated and eventually released to farmers as they indicated promising relationship with yield potential compared to other crosses. Crosses MUL 508 x MUL 516, POP A x MUL 141 and POP A x MUL 688 can further be evaluated for disease resistance. The results will be useful to breeders and farmers in selecting the potential parental materials for improvement in maize breeding programs.

Keywords: Combining ability; inbred line; positive; negative; estimates; combiner.

1. INTRODUCTION

Maize (*Zea mays*) also known as corn is an economically important crop worldwide and particularly in Africa where the annual production exceeds that of wheat and other important staple crops. Combining ability is of special importance in cross pollinated crops like maize because it helps in identifying potential inbred parents with hybrid vigor. Utilization of heterosis for genetic improvement of different traits and combining ability are fundamental tools for enhancing productivity in the form of filial one hybrid [1]. Farmers require high yielding maize hybrids with stable grain yield across environments that differ in terms of biotic and abiotic stress factors.

Heterosis increases yield potential and improves adaptation to stress in maize, however the underlying mechanism of heterosis and combining ability remains elusive [2]. A wide range of natural genetic diversity has been captured in the current maize germplasm [3,4].

The concept of general combining ability and specific combining ability (GCA & SCA) was introduced by Sprague and Tatum [5] and its mathematical modeling was set about by Griffing [6] in his classical paper in conjunction with the diallel crosses. The value of any population depends on its potential and its combining ability in crosses [7]. The average performance of a particular inbred in a series of hybrid combinations is known as its general combining ability while specific combining ability refers to the performance of a combination of two specific inbred lines in a particular cross [8]. Maize displays an orderly sequence of development of yield component namely; number of cobs per

plant, number of rows per cob, number of kernels per row and kernel weight [9,10,11,12,13] reported that GCA and SCA effects were highly significant but GCA effects on 100 grain weight under high temperature condition was non-significant. The use of diallel crosses to study genetic control of trait and to select parents to obtain synthetics or hybrids is frequent in maize breeding [14,15,16]. This study was conducted to evaluate the effects of GCA and SCA of the inbred lines using complete diallel scheme. The results in this study will be useful to maize breeders and farmers in selecting the potential parental materials for improvement in maize breeding programs in Kenya.

2. MATERIALS AND METHODS

2.1 Study Site

Field trials were grown at the KALRO experimental stations in Kiambu and Embu counties, during the long rain growing season from March to September 2012.

2.2 Study Materials

The germplasm used in the study were 18 inbred lines and their respective single crosses derived from KALRO, Muguga. The inbred lines were white, tropical lines and were rain fed throughout the growing season. The sets were without reciprocals (Table 1).

2.3 Experimental Layout

The experiment was laid out in a 6 x 6 lattice design consisting of 6 blocks with two replication

Mean separation was done using Tukey's comparison method at 0.05% significance level. Combining ability was estimated using Griffing (1956) analysis method, model 1 expressed as:

$$X_{ij} = u + g_i + g_j + s_{ij} + e$$

where:

u was the overall mean of all entries in the diallel design

g_i is the general combining ability of the i th parent

g_j was the general combining ability of the j th parent

s_{ij} was the specific combining ability between the i th and j th parent

e was the error.

3. RESULTS

3.1 Mean Performance on Different Morphological Traits in KALRO Embu

The analysis of variance (ANOVA), showing the mean squares of plant height (PH), ear height (EH), disease scores of maize streak virus (MSV) disease, grey leaf spot (GLS), and grain yield (GY) for Embu are shown in Table 2. The crosses showed a highly significant difference ($p < 0.001$) for plant height and ear height. They also showed a significant difference ($p < 0.05$) on grain yield.

3.2 Mean Performance in KALRO Muguga

The analysis of variance (ANOVA) for plant height, ear height, disease scores for MSV and GLS and grain yield for Muguga shown in Table 3 showed significant ($p < 0.05$) difference for plant height, ear height and GLS, while there was no significant difference on MSV and grain yield on the crosses in Muguga (Table 3).

General combining ability analysis revealed significant difference on plant height, ear height and grain yield. Data on GCA (Table 4) revealed that set one where MUL 508 was used the GCA range on grain yield was -0.8 to 0.5 t/ha, when POP A was used the GCA range on grain yield was -0.09 to 0.21 t/ha, while when inbred line MUL 513 was used the GCA range on grain yield was -0.18 to 0.62 t/ha. Data (Table 4) showed that inbred line MUL 508 had positive significant ($p < 0.05$) GCA for ear height when crossed with inbred line MUL 541, it also showed a positive significant GCA for grain yield when crossed with inbred line MUL 688. Inbred line MUL508 further showed a positive significant GCA for GLS disease resistance when crossed with inbred line CN 244 (Table 4). Data for set two showed inbred line POP A had a negative GCA -1.71 and -1.57 for plant height and ear height respectively when crossed with MUL 511, however there was positive GCA with grain yield 0.21, MSV disease resistance 0.31 and GLS disease resistance 0.71 (Table 4). The data further showed that when POP A was

Table 2. Performance of for different morphological traits in KALRO Embu

Source of variation	Df	Mean sum of squares				
		PH (cm)	EH(cm)	MSV	GLS	GY(t/ha)
Replication	1	6290.7	660.1	10.889	27.5035	3.19
Genotype	35	2047.7**	1221.8**	2.2	0.5527	4.02*
Error	35	175.8	169	1.203	0.5035	1.74
Overall mean		185	99.5	1.00	1.44	2.68
CV%		7.1	4.3	5.5	49.4	7.2

*,** Significant at ($p < 0.05$), and ($p < 0.001$) respectively, PH-plant height, EH-ear height, MSV-maize streak virus, GLS-grey leaf spot, GY-grain yield, CV%-Coefficient of variation

Table 3. Performance for different morphological traits in Muguga

SV	Df	Mean sum of squares				
		PH(cm)	EH(cm)	MSV	GLS	GY(t/ha)
Replication	1	19	32	0.0868	0	1.488
Genotype	35	2175.9**	796.8**	0.1725	0.2865**	11.536
Error	35	362.4	115.7	0.2225	0	8.215
Overall mean		214	79.9	1.16	1.94	4.84
CV%		0.3	1.2	4.2	0	4.2

** Significant at ($p < 0.05$), PH- Plant height, EH- Ear height, MSV-Maize streak virus, GLS- Grey leaf spot, GY- Grain yield, CV%-Coefficient of variation

Table 4. Estimates of G.C.A effects for inbred lines for yield and yield components for Embu and Muguga sites combined

Sets	Inbred lines	PH(CM)	EH(CM)	GY(T/HA)	Msvscores	GLS scores
MUL	MUL 516	34.17	19	-0.8	0.42	1.24
508	MUL 521	32.17	16	0.5	0.28	1.52
	MUL 141	-17.83	-14	0.1	-0.12	-0.36
	MUL 541	6.17	0.0**	0.5	-0.12	0.61
	MUL 688	57.93	-26	0.0**	0.08	-2.99
	CN 244	3.17	0.5	-0.3	0.28	0.01*
POP A	MUL 511	-1.71	-1.57	0.21	0.31	0.71
	MUL 521	-13.71	2.43	-0.09	-0.29	-0.34
	MUL 114	38.29	29.43	-0.29	-0.19	1.22
	MUL 141	-4.71	-14.57	-0.09	0.11	-0.31
	MUL 536	-12.71	-6.57	-0.19	-0.09	-0.29
	MUL 541	-24.71	-14.57	0.21	0.31	-2.62
	MUL 688	19.29	5.43	0.21	-0.19	1.66
MUL	MUL 531	13	7.8	0.18	-0.28	1.16
513	MUL 533	19	17.8	0.38	0.28	0.91
	MUL 114	-29	-23.2	-0.12	0.28	-1.3
	CN 244	-25	-12.2	0.62	-0.02*	-0.67
	MUL 516	22	9.8	-0.18	-0.32	-0.06

*, ** Significant at $p < 0.05$ and $p < 0.001$ levels respectively,

PH=Plant height, EH= Ear height, GY=Grain yield, MSV=Maize streak virus disease, GLS=Grey leaf spot

crossed with inbred line MUL 521 there was negative GCA in all the traits except ear height which had positive GCA of 2.43 cm. Further negative GCA was seen when the inbred line POP A was crossed with inbred line MUL 536 in all the traits studied, however with inbred line MUL 688 there was positive GCA for all traits

except for MSV disease resistance (-0.19). When inbred line MUL 513 in set three was crossed with inbred line MUL 531 there was a negative GCA for MSV resistance (-0.28) but a positive GCA for all the other traits. The data also showed positive GCA for all the traits when POP A was crossed with MUL 533 (Table 4).

Table 5. Estimates of S.C.A effects for inbred lines for yield and yield components for Embu and Muguga sites combined

Sets	Crosses	PH(CM)	EH(CM)	GY(T/ha)	MSV(Scores)
1	MUL508XMUL516	-8.33	-4.75	-0.31	0.16
	MUL508XMUL521	-8.08	-4.00	-0.38	-0.16
	MUL508XMUL141	4.42	3.50	0.09	0.06
	MUL508XMUL541	-1.58	0.0**	-0.15	-0.14
	MUL508XMUL688	14.42	6.5	0.75	0.04*
	MUL508XCN244	-0.83	-1.25	0.0**	0.04*
2	POPAXMUL511	-0.31	0.17	-0.18	-0.06
	POPAXMUL521	2.09	-0.63	0.03*	-0.06
	POPAXMUL114	-8.31	-6.03	-0.28	0.12
	POPAXMUL141	0.29	2.77	0.03*	0.14
	POPAXMUL536	1.89	1.17	0.02*	0.00**
	POPAXMUL541	4.29	2.77	0.49	-0.06
	POPAXMUL688	0.09	-0.23	-0.12	-0.06
3	MUL513XMUL531	-3.6	-1.93	-0.32	-0.08
	MUL513XMUL533	-5.6	-5.26	-0.23	-0.15
	MUL513XMUL114	10.4	8.40	0.51	0.02*
	MUL513XMUL516	9.1	4.74	0.28	0.19
	MUL513XCN244	-10.3	-5.93	-0.22	0.02*

*, ** Significant at $p < 0.001$ and $p < 0.05$ respectively

PH=Plant height (cm), EH=Ear height (cm), GY=Grain yield (t/ha), MSV=Maize Streak Virus, GLS=Grey Leaf Spot

Data on specific combining ability (SCA) (Table 5) showed that there was a significant ($p < 0.05$) difference between the inbred lines in the three sets on grain yield. The SCA range on grain yield in set one where MUL 508 was used was -0.15 to 0.75 t/ha, when POPA was used the SCA range on grain yield was -0.28 to 0.49 t/ha while for inbred line MUL 513 the SCA range was -0.32 to 0.51 (Table 5). Data on grain yield in set one showed a positive SCA on cross MUL 508 x MUL 141 (0.09) and MUL 508 x MUL 688 (0.75) (Table 5). Data further showed a positive SCA in set two on cross POP A x MUL 521 (0.03), POP A x MUL 141 (0.03), POP A x MUL 536 (0.02) and POP A x MUL 541 (0.49) on grain yield (Table 5). Further the data showed a positive SCA on yield on crosses MUL 513 x MUL 114 (0.05) and MUL 513 x MUL 516 (0.28).

3.3 Performance of Crosses on Both Research Environments Embu and Muguga

Data comparing the mean plant height for the two environments Embu and Muguga revealed that there was a significant ($P < 0.001$) difference on site and also on the crosses performance (Table 6).

Data on mean ear height in the two environments showed that there was a significant $P < 0.001$ difference between the site and crosses performance (Table 7).

Data on MSV disease score showed a significant $P < 0.001$ difference on the crosses (Table 8), however there was no significant $P > 0.05$ difference on the sites for MSV disease score.

Data on mean grey leaf spot scores showed there was significant ($P \leq 0.05$) difference among genotypes in the two research sites Embu and Muguga (Table 9).

Data on grain yield for the sampled crosses in both environments Embu and Muguga showed

there was significant ($P < 0.05$) difference in grain yield (t/ha) between the research sites (Table 10).

4. DISCUSSION

Data on combining ability where set 1, 2 and 3 (MUL 508, POP A and MUL 513) were crossed in diallel system showed the significant GCA and SCA variances for majority of the traits. This indicated the importance of both additive and non-additive gene actions in the expression of these traits. Diallel mating systems have provided genetic understanding for a chosen set of parents [17] and have been used to study various traits in many crops. Positive GCA deviation for yield indicated high yield potential as seen in cross MUL 508 x MUL 521 (0.5) and POP A x MUL 688 (0.21) (Table 5), while positive GCA values for disease scores indicated vulnerability to disease (Table 5). These observations were in line with those of [18], who observed that GCA and SCA variances were highly significant for yield and yield attributing traits which indicated the importance of both additive and non-additive gene action.

There was a significant difference among GCA and SCA effects on ear height but this trait was highly influenced by the additive effect since GCA contributed highest percentage to the crosses sum squares. MUL 508 showed significant ($p < 0.05$) difference when crossed with MUL 541 for ear height and grain yield. Most of the SCA estimates were non-significant, further indicating that performance of crosses for ear height was dependent on GCA effects (Table 5). Predominance of additive genetic effects in influencing ear height has been reported previously [19,20]. Inbred line MUL 508 showed significant ($P < 0.05$) difference when crossed with MUL 141, MUL 541, MUL 688 and CN 244 for GLS disease. MUL 508 also showed a highly significant ($p < 0.001$) difference when crossed with CN 244 for GLS disease scores.

Table 6. Performance of crosses on plant height in KARLO Embu and KALRO Muguga environments

Source of variation	d.f.	s.s.	m.s.	v.r.	F pr.
Rep	1	2809.0	2809.0	8.9	
Env	1	30044.4	30044.4	95.5	<.001
Genotype	35	131972.6	3770.6	12.0	<.001
Env/Genotype	35	15852.6	452.9	1.4	0.097
GCA	17	116.35	58.175	70.175	
SCA	35	0.05*	0.00361*	207.5	
Residual	123	22336.0	314.6		

Table 7. Performance for ear height in Embu and Muguga environments

Source of variation	d.f.	s.s.	m.s.	v.r.	F pr.
Rep	1	200.7	200.7	1.4	
Env.	1	13806.3	13806.3	93.8	<.001
Crosses	35	63608.8	1817.4	12.3	<.001
Env/Crosses	35	7041.2	201.2	1.4	0.133
GCA	17	4.49	2.245	200.265	
SCA	35	0.01*	0.000278*	207.01	
Residual	123	10456.3	147.3		

Table 8. Performance of crosses for MSV disease scores for Embu and Muguga environments

Source of variation	d.f.	s.s.	m.s.	v.r.	F pr.
Rep	1	0.16	0.16	0.03	
Env.	1	17.57	17.57	3.53	0.064
Crosses	35	377.41	10.78	2.17	0.003*
Env/Crosses	35	167.04	4.77	0.96	0.542
GCA	17	0.75	0.365	207.75	
SCA	35	0.1	0.000556*	207	
Residual	123	352.96	4.97		

Table 9. Performance on GLS scores at Embu and Muguga research sites

Source of variation	d.f.	s.s.	m.s.	v.r.	F pr.
Rep	1	4.516	4.516	5.690	
Env.	1	0.918	0.918	1.160	<.001
Crosses	35	37.769	1.079	1.360	0.286
Env/Crosses	35	45.269	1.293	1.630	0.041*
GCA	17	0.09	0.05*	207.09	
Residual	88	56.359	0.794		

Table 10. Performance of crosses for grain yield at Embu and Muguga research sites

Source of variation	d.f.	s.s.	m.s.	v.r.	F pr.
Rep	1	13.14	13.14	27.86	
Env.	1	15.02	15.02	31.84	<.001
Crosses	35	9.55	0.27	0.58	0.961
Env/Crosses	35	11.67	0.33	0.71	0.869
GCA	17	0.85	0.425	207.85	
SCA	35	0.07	0.000278	207	
Residual	123	33.48	0.47		

5. CONCLUSIONS AND RECOMMENDATIONS

GCA effects showed that inbred line MUL 513 was a good general combiner for grain yield, ear height and plant height. Inbred line POPA was a good general combiner for grain yield. Based on SCA estimates, the best cross combinations for plant height were MUL 508 x MUL 688, POP A x MUL 541 and MUL 513 x MUL 114. For ear height best cross combinations were: MUL 508 x MUL 688, POP A x MUL 141, POP A x MUL 541 and MUL 513 x MUL 114. The best combinations for grain yield were MUL 508 x MUL 688, POPA x MUL 541 and MUL 513 x MUL 114. Inbred line

MUL 513 showed positive GCA for plant height, ear height, and grain yield and a negative GCA for MSV when crossed with inbred line MUL 531, MUL 533 and CN244, showing additive genes were more predominant. These traits can be chosen as superior characteristics to help improve maize grain yield. Inbred lines MUL 508, POP A and MUL 513 can be used in improving other genotypes on disease resistant trait.

ACKNOWLEDGEMENTS

We are grateful to KALRO Muguga South and Embu centres for providing all the resources needed in the project.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Flint-Garcia SA, Buckler ES, Tiffin P, Ersoz E, Springer NM. Heterosis is prevalent for multiple traits in diverse maize germplasm. *Plos One*. 2009;4:e7433. 10 Journal.
2. Ararus J, Sanches C, Cabrera L. Is heterosis in maize mediated through better water use. *New Phytologist*. 2010;187:392-406.
3. Wright SI, Schroeder SG, Yamasaki M, Doebley JF, McMullen MD, Gaut BS. The effects of artificial selection on the maize genome. *Science*. 2005;308:1310-1314.
4. Troyer AF. Adaptedness and heterosis in corn mule hybrids. *Crop Science*. 2006;46:528-543.
5. Sprague GF, Tatum LA. General versus specific combining ability in single crosses of corn. *Journal of the American Society of Agronomy*. 1942;34:923-932.
6. Griffing B. Concept of general and specific combining ability in relation to diallel crossing systems. *Australian Journal of Biological Science*. 1956;9:463-493.
7. Vacaro E, Neto B, Fernards J, Pegoraro DJ, Nuss CN, Conceino LD. Combining ability of twelve maize populations. *Brazzillian Society of Genetics*. 2002;37:67-72.
8. Sharief AE, Elkalla SE, Gado HE, Abo-Yousef H. Heterosis in yellow maize. *Australian Journal of Crop Science*. 2009;3:146-154.
9. Viola G, Ganesh M, Reddy SS, Kummar CVS. Study of heritability and genetic advances in elite baby corn (*Zea mays*) lines. *Progressive Agriculture*. 2003;3(2):127-128. Introduction to Regression Test Automation. Available: [Software.www.operativesoft.com](http://www.operativesoft.com)
10. Kalla V, Kumar R, Basandrai AK. Combining ability and gene action estimates of yield and yield contributing characters in maize. *Crop Research*. 2001;22:102-106.
11. Renugopal MN, Ansar A, Rao NV. Combining ability studies in maize (*Zea mays* L.) Annual Agriculture Research. 2002;23(1):92-95.
12. Unay A, Konak C. Inheritance of grain yield in a half – diallel maize population. *Turkish Journal of Agriculture*. 2004;28:239-244.
13. Akbar M, Saleem M, Azhar FM, Yasin M, Ahmad R. Combining ability analysis in maize under normal and high temperature conditions. *Journal of Agricultural Research*. 2008;46(1):27-38.
14. Welcker C, The C, Andreau B, De Leon C, Parentoni SN, Bernal J, et al. Heterosis and combining ability for maize adaptation to tropical acid soils: Implications for future breeding strategies. *Crop Science*. 2005;45:2405-2413.
15. Rodrigues MC, Chaves LJ, Pacheco CAP. Heterosis in crosses among white grain, maize populations with high quality protein. *Brazzillian Society of Genetics*. 2006;41:59-66.
16. Vivek S, Odongo O, Njuguna J, Imanywona J, Diallo A, Pixley K. Diallel analyses of grain yield and resistance to seven diseases of 12 African maize (*Zea mays* L.) inbred lines. *Euphytica*. 2010;172:329-340.
17. Murray LW, Ray IM, Dong H, Segovialerma A. The gardener and eberhart analyses 11 and 111 re-visited. *Crop Science*. 2003;43:1930-1937.
18. Kumar S, Singh HB, Sharma JK. Combining ability analysis for grain yield and other associated traits in rice. *Oryza*. 2007;44(2):108-114.
19. Glover MA, Willmot DB, Darrah LL, Hibbard BE, Zhu X. Diallel analyses of agronomic traits using Chinese and U.S. maize germplasm. *Crop Science*. 2005;45:1096-1102.
20. Menkir A, Ayodele M. Genetic analysis of resistance to GLS of mid altitude maize inbred lines. *Crop Science*. 2005;45:163-170.

© 2016 Kariuki et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/15981>